arXiv:0710.5064v1 [gr-qc] 26 Oct 2007

Dirac quasinormal modes of a Schwarzschild black hole surrounded by free static spherically symmetric quintessence

Yu Zhang,[*] Yuan-Xing Gui,[†] and Fei Yu
School of Physics and Optoelectronic Technology,
Dalian University of Technology, Dalian, 116024, P. R. China

We evaluate the quasinormal modes of massless Dirac perturbation in a Schwarzschild black hole surrounded by the free static spherically symmetric quintessence by using the third-order WKB approximation. The result shows that due to the presence of quintessence, the massless field damps more slowly. The real part of the quasinormal modes increases and the the absolute value of the imaginary part increases when the state parameter w_q increases. In other words, the massless Dirac field decays more rapidly for the larger w_q. And the peak value of potential barrier gets higher as $|k|$ increases and the location of peak moves along the right for fixed w_q.

PACS numbers: 04.30.Nk, 04.70.Bw, 97.60.Lf
Keywords: Quasinormal modes; Dirac perturbation; WKB approximation.

I. INTRODUCTION

Quasinormal modes of black holes play a important role in the black hole theory. They are regards as the "characteristic sound"of black holes. The concept of QNMs was firstly pointed out by Vishveshwara[1] in calculations of the scattering of gravitational waves by a black hole. In recent decades, many people have done a lot of work of the quasinormal modes of black holes[2]-[10]. They have investigate the perturbation such as: gravitational perturbation[11], scalar perturbation[12], electromagnetic perturbation[13], Dirac perturbation[14]-[17], and so on. There are many methods to calculate the QNMs of the perturbation, for example, WKB approximation[18]-[20], the "potential fit"[21], and the method of continued fractions[22], etc. In this paper, we use the third WKB approximation.

There are three major motivations to study the QNMs in detail. First, QNMs may provide valuable help to identify black hole parameters[23][24]. Second, an important aspect of QNMs studies has been related to the AdS/CFT conjecture. By computing QN frequencies in AdS spcectime, we can obtain a prediction for the thermalization timescale in the strongly coupled CFT. It is argued that string theory in AdS space is equivalent to CFT in one less dimension[25]. And the third is the relation between QNMs and black hole area quantization[26]-[28].

Recently, V. V. Kiselev[29] has considered Einstein's field equations for a black hole surrounded by the static spherically symmetric quintessential matter and obtained a new solution that depends on the state parameter w_q of the quintessence. And Chen and Jing evaluated the quasinormal frequencies of massless scalar field perturbation[30] around the black hole which is surrounded by quintessence. In other papers, we have investigate the gravitational perturbation[31] and the electromagnetic perturbation[32]. In this paper, we consider the massless Dirac field in this situation.

II. DIRAC EQUATION IN THE SPACE TIME OF SCHWARZSCHILD BLACK HOLE SURROUNDED BY QUINTESSENCE

The metric for the spacetime of Schwarzschild black hole surrounded by the static spherically-symmetric quintessence is given by[30]

$$ds^2 = -fdt^2 + f^{-1}dr^2 + r^2(d\theta^2 + sin^2\theta d\phi^2), \tag{1}$$

with

$$f = 1 - \frac{2M}{r} - \frac{c}{r^{3w_q+1}} \tag{2}$$

---

[*] Electronic address: zhangyu128@student.dlut.edu.cn
[†] Electronic address: guiyx@dlut.edu.cn

where M is the black hole mass, w_q is the quintessential state parameter, c is the normalization factor related to $\rho_q = -\frac{c}{2}\frac{3w_q}{r^{3(1+w_q)}}$, and ρ_q is the density of quitenssence.
The massless Dirac equation in the black hole spacetime can be written as[33]

$$[\gamma^a e_a^\mu(\partial_\mu + \Gamma_\mu)]\Psi = 0, \tag{3}$$

where e_a^μ is the inverse of the tetrad e_μ^a defined by the metric $g_{\mu\nu}$,

$$g_{\mu\nu} = \eta_{ab} {e_\mu}^a {e_\nu}^b \tag{4}$$

with $\eta_{ab} = \mathrm{diag}(-1, 1, 1, 1)$ being the Minkowski metric.
γ^a is the Dirac matrix, and Γ_μ is the spin connection which is given by

$$\Gamma_\mu = \frac{1}{8}[\gamma^a, \gamma^b] e_a^\nu e_{b\nu;\mu} \tag{5}$$

where $e_{b\nu;\mu} = \partial_\mu e_{b\nu} - \Gamma^\alpha_{\mu\nu} e_{b\alpha}$ is the covariant derivative of $e_{b\nu}$, and $\Gamma^\alpha_{\mu\nu}$ is the Christoffel symbols.
we take the tetrad as

$$e_\mu^a = diag(\sqrt{f}, \frac{1}{\sqrt{f}}, r, r\sin\theta). \tag{6}$$

Γ_μ can be expressed as

$$\Gamma_0 = \frac{1}{4} f' \gamma^1\gamma^0, \Gamma_1 = 0, \Gamma_2 = \frac{1}{2}\sqrt{f}\gamma^1\gamma^2, \Gamma_3 = \frac{1}{2}(\sin\theta\sqrt{f}\gamma^1\gamma^3 + \cos\theta\gamma^2\gamma^3) \tag{7}$$

Then, the Dirac equation (3) becomes

$$\frac{\gamma^0}{\sqrt{f}}\frac{\partial\Psi}{\partial t} + \sqrt{f}\gamma^1\left(\frac{\partial}{\partial r} + \frac{1}{r} + \frac{1}{4f}\frac{df}{dr}\right)\Psi + \frac{\gamma^2}{r}(\frac{\partial}{\partial\theta} + \frac{1}{2}cot\theta)\Psi + \frac{\gamma^3}{r\sin\theta}\frac{\partial\Psi}{\partial\varphi} = 0. \tag{8}$$

Define

$$\Psi = f^{-\frac{1}{4}}\Phi, \tag{9}$$

Eq. (8) becomes

$$\frac{\gamma^0}{\sqrt{f}}\frac{\partial\Phi}{\partial t} + \sqrt{f}\gamma^1\left(\frac{\partial}{\partial r} + \frac{1}{r}\right)\Phi + \frac{\gamma^2}{r}(\frac{\partial}{\partial\theta} + \frac{1}{2}cot\theta)\Phi + \frac{\gamma^3}{r\sin\theta}\frac{\partial\Phi}{\partial\varphi} = 0. \tag{10}$$

Define the tortoise coordinate r_* as

$$r_* = \int \frac{dr}{f} \tag{11}$$

Introducing the ansatz

$$\Phi = \begin{pmatrix} \frac{iG^{(\pm)}(r)}{r}\phi^{\pm}_{jm}(\theta,\varphi) \\ \frac{F^{(\pm)}(r)}{r}\phi^{\mp}_{jm}(\theta,\varphi) \end{pmatrix} e^{-i\omega t}, \tag{12}$$

with spinor angular harmonics

$$\phi^+_{jm} = \begin{pmatrix} \sqrt{\frac{l+\frac{1}{2}+m}{2l+1}} Y_l^{m-1/2} \\ \sqrt{\frac{l+\frac{1}{2}-m}{2l+1}} Y_l^{m+1/2} \end{pmatrix}, \qquad (for \quad j = l + \frac{1}{2}),$$

$$\phi^-_{jm} = \begin{pmatrix} \sqrt{\frac{l+\frac{1}{2}-m}{2l-1}} Y_l^{m-1/2} \\ -\sqrt{\frac{l+\frac{1}{2}+m}{2l-1}} Y_l^{m+1/2} \end{pmatrix}, \qquad (for \quad j = l - \frac{1}{2}),$$

equations 10 can be written in the form

$$\begin{pmatrix} 0 & -\omega \\ \omega & 0 \end{pmatrix} \begin{pmatrix} F^{\pm} \\ G^{\pm} \end{pmatrix} - \frac{\partial}{\partial r_*} \begin{pmatrix} F^{\pm} \\ G^{\pm} \end{pmatrix} + \sqrt{f} \begin{pmatrix} \frac{k_{\pm}}{r} & 0 \\ 0 & -\frac{k_{\pm}}{r} \end{pmatrix} \begin{pmatrix} F^{\pm} \\ G^{\pm} \end{pmatrix} = 0. \tag{13}$$

The cases for (+) and (−) in the functions $F^{\pm}$ and $G^{\pm}$ can be put together giving

$$\frac{d^2F}{dr_*^2} + (\omega^2 - V_1)F = 0, \tag{14}$$

$$\frac{d^2G}{dr_*^2} + (\omega^2 - V_2)G = 0, \tag{15}$$

with

$$V_1 = \frac{\sqrt{f}|k|}{r^2}\left(|k|\sqrt{f} + \frac{r}{2}\frac{df}{dr} - f\right), \quad \left(for \ \ k = j + \frac{1}{2}, \qquad and \ \ j = l + \frac{1}{2}\right), \tag{16}$$

$$V_2 = \frac{\sqrt{f}|k|}{r^2}\left(|k|\sqrt{f} - \frac{r}{2}\frac{df}{dr} + f\right), \quad \left(for \ \ k = -\left(j + \frac{1}{2}\right), \ \ and \ \ j = l - \frac{1}{2}\right). \tag{17}$$

The potentials V_1 and V_2 are supersymmetric partners derived from the same superpotential. We shall concentrate just on Eq. 14 with potential V_1 in evaluating the quasinormal mode frequencies for the massless Dirac field by the third WKB approximation in the following sections.

III. MASSLESS DIRAC QUASINORMAL FREQUENCIES

In this section we evaluate the QN frequencies for the massless Dirac field in the space time of the Schwarzschild black hole surrounded by quintessence using the third WKB approximation. WKB method was firstly developed by B. F. Schutz and C. M. Will[18] at the second order. later, S. Iyer and C. M. Will[19] developed the method to the third order and R. A. Konoplya[20] extended it to the sixth order.The accuracy of the WKB formula is better with a larger multipole number l and a smaller overtone n. The formula for the complex quasinormal frequencies ω(for the third order WKB method) is

$$\omega^2 = \left[V_0 + (-2V_0'')^{1/2}\Lambda\right] - i(n + \frac{1}{2})(-2V_0'')^{1/2}(1 + \Omega) \tag{18}$$

where

$$\Lambda = \frac{1}{(-2V_0'')^{1/2}}\left\{\frac{1}{8}\left(\frac{V_0^{(4)}}{V_0''}\right)(\frac{1}{4} + \alpha^2) - \frac{1}{288}\left(\frac{V_0'''}{V_0''}\right)^2(7 + 60\alpha^2)\right\}$$

$$\begin{aligned}\Omega = \frac{1}{-2V_0''}\Big\{&\frac{5}{6912}\left(\frac{V_0'''}{V_0''}\right)^4(77 + 188\alpha^2) \\ &- \frac{1}{384}\left(\frac{V_0'''^2V_0^{(4)}}{V_0''^3}\right)(51 + 100\alpha^2) + \frac{1}{2304}\left(\frac{V_0^{(4)}}{V_0''}\right)^2(67 + 68\alpha^2) \\ &+ \frac{1}{288}\left(\frac{V_0'''V_0^{(5)}}{V_0''^2}\right)(19 + 28\alpha^2) - \frac{1}{288}\left(\frac{V_0^{(6)}}{V_0''}\right)(5 + 4\alpha^2)\Big\}\end{aligned} \tag{19}$$

and

$$\alpha = n + \frac{1}{2}, \ V_0^{(n)} = \frac{d^nV}{dr_*^n}\Big|_{r_* = r_*(r_p)} \tag{20}$$

Take $M = 1, c = 0.001$ and $M = 1, c = 0$ for our calculation. And $c = 0$ means there is no quintessence. Using the third-order WKB approximation, we can get the solutions as the table 1 and table 2 show(for $0 \leqslant n < k$), where l is the angular harmonic index, n is the overtone number, ω is the complex quasinormal frequencies, w_q is the



FIG. 1: Variation of the effective potential for the massless Dirac field with $|k|$ in the Schwarzshild black hole surrounded by quintessence for $w_q = -1/2$ and $c = 0.001$.



FIG. 2: Variation of the effective potential for the massless Dirac field in the Schwarzshild black hole surrounded by quintessence with r for $|k| = 5, c = 0.001$ and $3w_q + 1 = -0.01, -1.7, -1.99$.

quintessential state parameter.

The effective potential $V(r, k)$, which depends on the absolute value of k and w_q, is in the form of a barrier. In Fig. 1, it shows the variation of effective potential with $|k|$ when $w_q = -\frac{1}{2}$ and $c = 0.001$. The variation of the effective potential with r which is respective to the quintessential state w_q parameter for fixed $|k| = 5$ and $c = 0.001$ is shown in Fig. 2.

TABlE I:The quasinormal frequencies of massless Dirac perturbations in the Schwarzshild black hole[14] without

quintessence(c=0).

$\lvert k\rvert$	n	ω	$\lvert k\rvert$	n	ω
1	0	$0.17645-0.10011i$	4	0	$0.76719-0.09628i$
				1	$0.75396-0.29105i$
				2	$0.73045-0.49088i$
				3	$0.69992-0.69571i$
2	0	$0.37863-0.09654i$	5	0	$0.96021-0.09626i$
	1	$0.35360-0.29875i$		1	$0.94959-0.29018i$
3	0	$0.57369-0.09632i$		2	$0.92998-0.48763i$
	1	$0.55619-0.29298i$		3	$0.90358-0.68924i$
	2	$0.52729-0.49719i$		4	$0.87205-0.89441i$

TABlE II: The quasinormal frequencies of massless Dirac perturbations in the black hole surrounded by quintessence for $|k|=2$, $|k|=3$, $|k|=4$, $|k|=5$ and $c=0.001$.

$3w_q+1$	$\omega(n=0)$	$3w_q+1$	$\omega(n=0)$	$\omega(n=1)$
$\lvert k\rvert=1$		$\lvert k\rvert=2$		
-0.01	$0.17620-0.09990i$	-0.01	$0.37806-0.09635i$	$0.35310-0.29813i$
-0.4	$0.17611-0.09977i$	-0.4	$0.37780-0.09623i$	$0.35287-0.29776i$
-0.8	$0.17598-0.09956i$	-0.8	$0.37741-0.09605i$	$0.35252-0.29721i$
-1.2	$0.17577-0.09926i$	-1.2	$0.37683-0.09579i$	$0.35199-0.29639i$
-1.6	$0.17546-0.09881i$	-1.6	$0.37596-0.09541i$	$0.35118-0.29523i$
-1.99	$0.17500-0.09819i$	-1.99	$0.37468-0.09489i$	$0.35001-0.29364i$

$3w_q+1$	$\omega(n=0)$	$\omega(n=1)$	$\omega(n=2)$	$\omega(n=3)$	$\omega(n=4)$
$\lvert k\rvert=3$					
-0.01	$0.57282-0.09613i$	$0.55536-0.29238i$	$0.52654-0.49617i$		
-0.4	$0.57240-0.09601i$	$0.55497-0.29203i$	$0.52618-0.49557i$		
-0.8	$0.57176-0.09584i$	$0.55436-0.29150i$	$0.52561-0.49466i$		
-1.2	$0.57079-0.09559i$	$0.55344-0.29073i$	$0.52476-0.49336i$		
-1.6	$0.56933-0.09524i$	$0.55205-0.28965i$	$0.52349-0.49152i$		
-1.99	$0.56718-0.09478i$	$0.55005-0.28822i$	$0.52169-0.48903i$		
$\lvert k\rvert=4$					
-0.01	$0.76603-0.09608i$	$0.75283-0.29046i$	$0.72938-0.48988i$	$0.69892-0.69429i$	
-0.4	$0.76546-0.09597i$	$0.75227-0.29011i$	$0.72885-0.48929i$	$0.69843-0.69345i$	
-0.8	$0.76457-0.09579i$	$0.75141-0.28959i$	$0.72803-0.48841i$	$0.69765-0.69220i$	
-1.2	$0.76321-0.09555i$	$0.75009-0.28884i$	$0.72678-0.48715i$	$0.69649-0.69042i$	
-1.6	$0.76116-0.09521i$	$0.74810-0.28782i$	$0.72491-0.48540i$	$0.69476-0.68791i$	
-1.99	$0.75812-0.09478i$	$0.74521-0.28649i$	$0.72224-0.48308i$	$0.69235-0.68452i$	
$\lvert k\rvert=5$					
-0.01	$0.95876-0.09606i$	$0.94817-0.28959i$	$0.92860-0.48664i$	$0.90226-0.68783i$	$0.87081-0.89258i$
-0.4	$0.95803-0.09595i$	$0.94745-0.28925i$	$0.92791-0.48606i$	$0.90160-0.68701i$	$0.87019-0.89151i$
-0.8	$0.95689-0.09578i$	$0.94633-0.28873i$	$0.92682-0.48519i$	$0.90057-0.68578i$	$0.86920-0.88992i$
-1.2	$0.95515-0.09554i$	$0.94462-0.28800i$	$0.92518-0.48396i$	$0.89899-0.68404i$	$0.86772-0.88766i$
-1.6	$0.95250-0.09521i$	$0.94203-0.28701i$	$0.92270-0.48227i$	$0.89666-0.68162i$	$0.86554-0.88449i$
-1.99	$0.94856-0.09480i$	$0.93823-0.28575i$	$0.91913-0.48009i$	$0.89336-0.67843i$	$0.86253-0.88020i$



FIG. 3: The relationship between the real and imaginary parts of quasinormal frequencies for the massless Dirac perturbations in the background of the black hole surrounded by quintessence for fixed $c = 0.001$ and no quintessence for $c = 0$. × and $*$, respectively, refering to the value with quintessence and without quintessence.

IV. DISCUSSION AND CONCLUSION

The quasinormal modes of a black hole present complex frequencies, the real part of which represents the actual frequencies of the oscillation and the imaginary part represents the damping.

From Fig.1 we see that the peak value of potential barrier gets higher as $|k|$ increases and the location of peak moves along the right for fixed w_q. From Fig. 2, it shows that when the absolute w_q increases, the peak value of potential barrier gets lower. Using the third WKB approximation, we calculate the lowly decaying modes frequencies for $k = 1, 2, 3, 4, 5$. The complex quasinormal mode frequencies for the massless Dirac field are listed in Table I(without quintessence) and Table II(exist quintessence, and c=0.001). We plot the relationship between the real and imaginary parts of quasinormal frequencies with the variation of w_q(for fixed $c = 0.001$), compared with the situation without quintessence. From Fig.3 we can find that for fixed c(unequal to 0) and l the absolute values of the real and imaginary parts decrease as the absolute value of the quintessence state parameter w_q increases. It means that when the absolute value of w_q is bigger, the oscillations damp more slowly. And the absolute values of the real and imaginary parts of quasinormal modes with quintessence are smaller compared with those with no quintessence for given l and n. That is to say, due to the presence of quintessence, the oscillations of the massless Dirac fields damp more slowly.

Acknowledgments

Yu Zhang wishes to thank Ph.D LiXin Xu for his helpful discussions. This work is supported by the National Natural Science Foundation of China under Grant No. 10573004.

---

[1] C. V. Vishveshwara 1970 Nature **227** 936
[2] K. D. Kokkotas and B. G. Schmidt 1999 Living Rev. Rel. **2** 2
[3] H. P. Nollert 1999 Class. Quant. Grav. **16** R159
[4] S. Chandrasekhar 1975 Proc. R. Soc. London, Ser. A. **343** 289-298
[5] V. Cardoso, J. P. S. Lemos, S. Yoshida 2004 Phys. Rev. D **70** 124032
[6] R. A. Konoplya 2005 Phys. Rev. D **71**, 024038
[7] E. Berti, Vitor Cardoso 2006 Phys. Rev. D **74** 104020
[8] S. B. Chen, B. Wang, R. K. Su 2006 Class. Quant. Grav. **23** 7581-7590
[9] R. A. Konoplya, A. Zhidenko 2006 Phys. Rev. D **73** 124040
[10] A. López-Ortega 2006 Gen.Rel.Grav. **38** 743-771
[11] M. Giammatteo and I. G. Moss 2005 Class. Quantum Grav. **22** 1803-1824
[12] R. A. Konoplya and A. V. Zhidenko 2006 Phys. Rev. D **73** 124040
[13] A. López-Ortega 2006 Gen.Rel.Grav. **38** 1747-1770
[14] H. T. Cho 2003 Phys. Rev. D **68** 024003
[15] M. Giammatteo and J. L. Jing 2005 Phys. Rev. D **71** 024007
[16] J. L. Jing 2004 Phys. Rev. D **69** 084009
[17] J. L. Jing and Q. Y. Pan 2005 Phys. Rev. D **71** 124011
[18] B. F. Schutz, and C. M. Will 1985 Astrophys. J. **291** L33-L36
[19] S. Iyer and C. M. Will 1987 Phys. Rev. D **35** 3621
[20] R. A. Konoplya 2003 Phys. Rev. D **68** 024018
[21] V. Ferrari and B. Mashhoon 1984 Phys. Rev. D **30** 295
[22] E. W. Leaver 1985 Proc. R. Soc. A **402** 285
[23] F. Echeverria 1989 Phys. Rev. D **40** 3194
[24] L. S. Finn 1992 Phys. Rev. D **46** 5236
[25] E. Witten 1998 Adv. Theor. Math. Phys. **2** 253
[26] J. Bekenstein 1974 Lett. Nuovo Cimento **11** 467
[27] J. Bekenstein 1998 in Cosmology and Gravitation ,edited by M. Novello (Atlasciences, France 2000) pp. 1-85, gr-qc/9808028
[28] S. Hod 1998 Phys. Rev. Lett. **81** 4293
[29] V. V. Kiselev 2003 Class. Quant. Grav. **20** 1187-1197
[30] S. B. Chen, J. L. Jing 2005 Class. Quant. Grav. **22** 4651-4657
[31] Yu Zhang, Y. X. Gui 2006 Class. Quantum Grav. **23** 6141-6147
[32] Yu Zhang, Yuanxing Gui et al 2007 Gen. Rel. Grav. **39** 1003-1010.
[33] D. R. Brill and J. A. Wheeler 1957 Rev. Mod. Phys. **29** 465